Ricin Fact Sheet



Castor beans (Source: U.S. Agricultural Research Service)

Ricin is a potent toxin that could be used as an agent of biological warfare or bioterrorism. It is a phytotoxalbumin protein derived from the beans of the castor plant (*Ricinus communis*). Ricin is widely available and easily produced. Ricin's significance as a potential bioweapon relates in part to its availability. Although ricin is not an ideal bioweapon agent, it remains a threat.

Delivery: Ricin can be delivered via inhalation, injection, or ingestion. Liquid or crystalline ricin could be utilized as a food or water contaminant. For ricin to be delivered as an aerosol, it must be lyophilized-a process by which a liquid substance is dried by freezing in a high vacuum. While it is not overly difficult to aerosolize ricin,

Quick Facts

Agent Type: Toxin Lethality: Moderate Transmission: None Treatment: None Status: Select Agent; CDC Category B Agent Delivery: Inhalation, injection, ingestion

Mechanism: Ricin prevents cells in the body from synthesizing proteins, which can lead to widespread organ damage as well as pulmonary, liver, renal, and immunological failure.

Effects: Fever, coughing, and gastrointestinal problems are likely to be the first symptoms. If eaten, the toxin causes stomach irritation, gastroenteritis, bloody diarrhea, and vomiting. Inhalation causes severe lung damage, including

pulmonary edema. It can

also cause



seizures and central nervous system depression. When inhaled, this toxin may produce pathologic changes within 8 hours and severe

Ricin protein structure (3-D ribbon drawing based on X-ray crystallography data*)

respiratory symptoms followed by acute respiratory failure in 36-72 hours. When ingested, ricin causes severe this step does present a technical challenge.

Production: Ricin can be produced

gastrointestinal symptoms followed by vascular collapse and death. Since ricin is a relatively large protein, it is not easily absorbed across the skin and dermal relatively easily and inexpensively in exposure to ricin is of little concern.

large quantities in a fairly lowtechnology setting. Worldwide, one million tons of castor beans are of castor oil; the waste mash from this process is 5-10% ricin by weight.

Treatment: There is no treatment or prophylaxis currently available for ricin processed annually in the production exposure. If exposure does not prove fatal within 3-5 days, the victim will usually recover.

Historic Use: Ricin has reportedly been used as a tool for assassinations. During the 1980's Iran-Iraq war, ricin may have been used. In February 2004, ricin was detected in mail received at the U.S. Senate office complex.

(Sources: Scotland Yard, CDC, U.S. Army, eMedicine.com)